

August, 2007  
OCU-PHYS 273

# Closed conformal Killing-Yano tensor and Kerr-NUT-de Sitter spacetime uniqueness

Tsuyoshi Houri<sup>a\*</sup>, Takeshi Oota<sup>b†</sup> and Yukinori Yasui<sup>a‡</sup>

<sup>a</sup> *Department of Mathematics and Physics, Graduate School of Science,  
Osaka City University  
3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, JAPAN*

<sup>b</sup> *Osaka City University Advanced Mathematical Institute (OCAMI)  
3-3-138 Sugimoto, Sumiyoshi, Osaka 558-8585, JAPAN*

## Abstract

We study spacetimes with a closed conformal Killing-Yano tensor. It is shown that the  $D$ -dimensional Kerr-NUT-de Sitter spacetime constructed by Chen-Lü-Pope is the only spacetime admitting a rank-2 closed conformal Killing-Yano tensor with a certain symmetry.

---

\* `hour@sci.osaka-cu.ac.jp`

† `toota@sci.osaka-cu.ac.jp`

‡ `yasui@sci.osaka-cu.ac.jp`

Higher dimensional black hole solutions have attracted renewed interests in the recent developments of supergravity and superstring theories. Recently, the  $D$ -dimensional Kerr-NUT-de Sitter metrics were constructed by [1]. All the known vacuum type D black hole solutions are included in these metrics [2]. Kerr-NUT-de Sitter metrics are also interesting from the point of view of AdS/CFT correspondence. Indeed, odd-dimensional metrics lead to Sasaki-Einstein metrics by taking BPS limit [1, 3, 4, 5] and even-dimensional metrics lead to Calabi-Yau metrics in the limit [1, 6, 7]. Especially, the five-dimensional Sasaki-Einstein metrics have emerged quite naturally in the AdS/CFT correspondence.

On the other hand, it has been shown that geodesic motion in the Kerr-NUT-de Sitter spacetime is integrable for all dimensions [8, 9, 10, 11, 12, 13]. Indeed, the constants of motion that are in involution can be explicitly constructed from a rank-2 closed conformal Killing-Yano (CKY) tensor. In this paper, using a geometric characterisation of the separation of variables in the Hamilton-Jacobi equation [14], we study spacetimes with a rank-2 closed CKY tensor.

The rank-2 CKY tensor is defined as a two-form

$$h = \frac{1}{2} h_{ab} dx^a \wedge dx^b, \quad h_{ab} = -h_{ba} \quad (1)$$

satisfying the equation [15]

$$\nabla_a h_{bc} + \nabla_b h_{ac} = 2\xi_c g_{ab} - \xi_a g_{bc} - \xi_b g_{ac}. \quad (2)$$

The vector field  $\xi_a$  is called the associated vector of  $h_{ab}$ , which is given by

$$\xi_a = \frac{1}{D-1} \nabla^b h_{ba}. \quad (3)$$

**Theorem** *Let us assume the existence of a single rank-2 CKY tensor  $h$  for  $D$ -dimensional spacetime  $(M, g)$  satisfying the conditions,*

$$(a) \, dh = 0, \quad (b) \, \mathcal{L}_\xi g = 0, \quad (c) \, \mathcal{L}_\xi h = 0. \quad (4)$$

*Then,  $M$  is only the Kerr-NUT-de Sitter spacetime<sup>1</sup>.*

The Kerr-NUT-de Sitter metric takes the form [1]:

(a)  $D = 2n$

$$g = \sum_{\mu=1}^n \frac{dx_\mu^2}{Q_\mu} + \sum_{\mu=1}^n Q_\mu \left( \sum_{j=0}^{n-1} A_\mu^{(j)} d\psi^j \right)^2. \quad (5)$$

---

<sup>1</sup>We require a further technical condition which will be detailed in the proof. See the assumption below eq.(14).

(b)  $D = 2n + 1$

$$g = \sum_{\mu=1}^n \frac{dx_\mu^2}{Q_\mu} + \sum_{\mu=1}^n Q_\mu \left( \sum_{j=0}^{n-1} A_\mu^{(j)} d\psi^j \right)^2 + S \left( \sum_{j=0}^n A^{(j)} d\psi^j \right)^2. \quad (6)$$

The functions  $Q_\mu$  are given by

$$Q_\mu = \frac{X_\mu}{U_\mu}, \quad U_\mu = \prod_{\substack{\nu=1 \\ (\nu \neq \mu)}}^n (x_\mu^2 - x_\nu^2), \quad (7)$$

where  $X_\mu$  is an arbitrary function depending only on  $x_\mu^2$  and

$$A_\mu^{(k)} = \sum_{\substack{1 \leq \nu_1 < \dots < \nu_k \leq n \\ (\nu_i \neq \mu)}} x_{\nu_1}^2 x_{\nu_2}^2 \dots x_{\nu_k}^2, \quad A^{(k)} = \sum_{1 \leq \nu_1 < \dots < \nu_k \leq n} x_{\nu_1}^2 x_{\nu_2}^2 \dots x_{\nu_k}^2, \quad (8)$$

( $A_\mu^{(0)} = A^{(0)} = 1$ ) and  $S = c/A^{(n)}$  with a constant  $c$ .

In the following we briefly describe the proof (see [16] for detailed analysis). The wedge product of two CKY tensors is again a CKY tensor and so the wedge powers  $h^{(j)} = h \wedge \dots \wedge h$  are CKY tensors. The condition (a) means that the Hodge dual  $(D - 2j)$ -forms  $f^{(j)} = *h^{(j)}$  are Killing-Yano tensors:

$$\nabla_{(a_1} f_{a_2) a_3 \dots a_{D-2j+1}}^{(j)} = 0. \quad (9)$$

These Killing-Yano tensors generate the rank-2 Killing tensors  $K^{(j)}$  obeying the equation  $\nabla_{(a} K_{bc)}^{(j)} = 0$ . Under the condition (a) the Killing tensors  $K^{(j)}$  are mutually commuting [12, 13],

$$[K^{(i)}, K^{(j)}]_S = 0. \quad (10)$$

The bracket  $[ \ , \ ]_S$  represents a symmetric Schouten product. The equation can be written as

$$K_{d(a}^{(i)} \nabla^d K_{bc)}^{(j)} - K_{d(a}^{(j)} \nabla^d K_{bc)}^{(i)} = 0. \quad (11)$$

Let us define the vector fields  $\eta^{(j)}$  by

$$\eta_a^{(j)} = K_a^{(j) b} \xi_b. \quad (12)$$

---

<sup>2</sup> We call the metric Kerr-NUT-de Sitter for an arbitrary  $X_\mu$ . The existence of  $h$  does not restrict the metric to be Einstein.

Then we have

$$\nabla_{(a}\eta_{b)}^{(j)} = \frac{1}{2}\mathcal{L}_\xi K_{ab}^{(j)} - \nabla_\xi K_{ab}^{(j)}, \quad (13)$$

which vanishes by the conditions (b) and (c), i.e.  $\eta^{(j)}$  are Killing vectors. We can show that Killing vectors  $\eta^{(i)}$  and Killing tensors  $K^{(j)}$  are mutually commuting [14],

$$[\eta^{(i)}, K^{(j)}]_S = 0, \quad [\eta^{(i)}, \eta^{(j)}] = 0. \quad (14)$$

Here, we assume that the Killing tensors  $K^{(j)}$  and  $K^{(ij)} = \eta^{(i)} \otimes \eta^{(j)} + \eta^{(j)} \otimes \eta^{(i)}$  are independent. Therefore all the separability conditions of the geodesic Hamilton-Jacobi equation are satisfied [14].

Let  $y^a$  be geodesic separable coordinates of  $D = (n + k)$ -dimensional spacetime  $M$ :

$$y^a = (x^\mu, \psi^i), \quad \mu = 1, 2, \dots, n, \quad i = 0, 1, \dots, k-1, \quad (15)$$

where  $k = n$  ( $k = n + 1$ ) for  $D$  even (odd). In these coordinates the commuting Killing vectors  $\eta^{(j)}$  ( $j = 0, 1, \dots, k-1$ ) are written as  $\eta^{(j)} = \partial/\partial\psi^j$ . From [17, 18, 19] the inverse metric components are of the form,

$$g^{\mu\mu} = \bar{\phi}_{(0)}^\mu(x), \quad g^{ij} = \sum_{\mu=1}^n \zeta_\mu^{ij}(x^\mu) \bar{\phi}_{(0)}^\mu(x), \quad (16)$$

and the components of the Killing tensors  $K^{(j)}$  are given by

$$K^{(j)\mu\nu} = \delta^{\mu\nu} \bar{\phi}_{(j)}^\mu(x), \quad K^{(j)\mu i} = 0, \quad K^{(j)i\ell} = \sum_{\mu=1}^n \zeta_\mu^{i\ell}(x^\mu) \bar{\phi}_{(j)}^\mu(x). \quad (17)$$

Here,  $\bar{\phi}_{(j)}^\mu$  is the  $j$ -th column of the inverse of an  $n \times n$  Stäckel matrix  $(\phi_\mu^{(j)})$ , i.e. each element depends on the variable corresponding to the lower index only:  $\phi_\mu^{(j)}(x^\mu)$ . It should be noticed that the Killing tensors are constructed from CKY tensors, so that they obey the following recursion relations as linear operators [14]:

$$K^{(j)} = A^{(j)}I - QK^{(j-1)}, \quad (18)$$

where  $I$  is an identity operator and  $Q$  is defined by

$$Q^a{}_b = -h^a{}_c h^c{}_b. \quad (19)$$

Here  $A^{(j)}$  is given by

$$\det^{1/2}(I + \beta Q) = \sum_{j=0}^n A^{(j)} \beta^j. \quad (20)$$

Note that the equation (2) with the condition (a) is equivalent to

$$\nabla_a h_{bc} = \xi_c g_{ab} - \xi_b g_{ac}. \quad (21)$$

We can further restrict the unknown functions  $\bar{\phi}_{(0)}^\mu$  and  $\zeta_\mu^{ij}$  in the metric (16). This is analyzed by considering the equation (21) with  $\xi = \eta^{(0)}$ , and finally we find the Kerr-NUT-de Sitter metric (5) or (6).

As a crosscheck of our theorem, we confirmed by the direct calculation that a CKY tensor satisfying (a), (b) and (c) does not exist in the five-dimensional black ring background [20].

## Acknowledgements

We would like to thank Yoshitake Hashimoto and Ryushi Goto for discussions. This work is supported by the 21 COE program “Construction of wide-angle mathematical basis focused on knots”. The work of YY is supported by the Grant-in Aid for Scientific Research (No. 19540304 and No. 19540098) from Japan Ministry of Education. The work of TO is supported by the Grant-in Aid for Scientific Research (No. 18540285 and No. 19540304) from Japan Ministry of Education.

## References

- [1] W. Chen, H. Lü and C.N. Pope, “General Kerr-NUT-AdS metrics in all dimensions,” *Class. Quant. Grav.* **23** (2006) 5323-5340, [arXiv:hep-th/0604125](#).
- [2] N. Hamamoto, T. Houri, T. Oota and Y. Yasui, “Kerr-NUT-de Sitter curvature in all dimensions,” *J. Phys.* **A40** (2007) F177-F184, [arXiv:hep-th/0611285](#).
- [3] Y. Hashimoto, M. Sakaguchi and Y. Yasui, “Sasaki-Einstein Twist of Kerr-AdS Black Holes,” *Phys. Lett.* **B 600**, 270 (2004), [arXiv:hep-th/0407114](#).
- [4] M. Cvetič, H. Lü, D.N. Page and C.N. Pope, “New Einstein-Sasaki spaces in five and higher dimensions,” *Phys. Rev. Lett.* **95**, 071101 (2005), [arXiv:hep-th/0504225](#).
- [5] M. Cvetič, H. Lü, D.N. Page and C.N. Pope, “New Einstein-Sasaki and Einstein spaces from Kerr-de Sitter,” [arXiv:hep-th/0505223](#).
- [6] T. Oota and Y. Yasui, “Explicit Toric Metric on Resolved Calabi-Yau Cone,” *Phys. Lett.* **B 639**, 54 (2006), [arXiv:hep-th/0605129](#).
- [7] H. Lü and C.N. Pope, “Resolutions of Cones over Einstein-Sasaki Spaces,” [arXiv:hep-th/0605222](#).
- [8] V.P. Frolov and D. Kubizňák, “‘Hidden’ Symmetries of Higher Dimensional Rotating Black Holes,” *Phys. Rev. Lett.* **98** (2007) 11101, [arXiv:gr-qc/0605058](#).
- [9] D. Kubizňák and V.P. Frolov, “Hidden Symmetry of Higher Dimensional Kerr-NUT-AdS Spacetimes,” *Class. Quant. Grav.* **24** (2007) F1-F6, [arXiv:gr-qc/0610144](#).
- [10] D.N. Page, D. Kubizňák, M. Vasudevan and P. Krtouš, “Complete Integrability of Geodesic Motion in General Kerr-NUT-AdS Spacetimes,” *Phys. Rev. Lett.* **98** (2007) 061102, [arXiv:hep-th/0611083](#).
- [11] V.P. Frolov, P. Krtouš and D. Kubizňák, “Separability of Hamilton-Jacobi and Klein-Gordon Equations in General Kerr-NUT-AdS Spacetimes,” *JHEP* **0702** (2007) 005, [arXiv:hep-th/0611245](#).
- [12] P. Krtouš, D. Kubizňák, D.N. Page and V.F. Frolov, “Killing-Yano Tensors, Rank-2 Killing Tensors, and Conserved Quantities in Higher Dimensions,” *JHEP* **0702** (2007) 004, [arXiv:hep-th/0612029](#).

- [13] P. Krtouš, D. Kubizňák, D.N. Page and M. Vasudevan, “Constants of Geodesic Motion in Higher-Dimensional Black-Hole Spacetime,” [arXiv:hep-th/0707.0001](#).
- [14] T. Houri, T. Oota and Y. Yasui, “Closed conformal Killing-Yano tensor and geodesic integrability,” [arXiv:hep-th/0707.4039](#).
- [15] S. Tachibana, “On conformal Killing tensor in a Riemannian space,” *Tôhoku Math. J.* **21** (1969) 56-64.
- [16] T. Houri, T. Oota and Y. Yasui, in preparation.
- [17] S. Benenti and M. Francaviglia, “Remarks on Certain Separability Structures and Their Applications to General Relativity,” *Gen. Rel. Grav.* **10** (1979) 79-92.
- [18] E.G. Kalnins and W. Miller, Jr., “Killing tensors and nonorthogonal variable separation for Hamiltonian-Jacobi equations,” *SIAM J. Math. Anal.* **12** (1981) 617-629.
- [19] S. Benenti, “Separation of Variables in the Geodesic Hamilton-Jacobi Equation,” in *Symplectic Geometry and Mathematical Physics*, Progress in Mathematics Vol. **99**, ed. by P. Donato, C. Duval, J. Elhadad and G.M. Tuynman, Birkhäuser, (1991) 1-36.
- [20] R. Emparan and H.S. Reall, “A rotating black ring in five dimensions,” *Phys. Rev. Lett.* **88** (2002) 101101, [arXiv:hep-th/0110260](#).